



ARL-TN-0749 • APR 2016



Theoretical Prediction of the Heats of Formation, Densities, and Relative Sensitivities for 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo (3:3:0)octan-2-one

by Edward FC Byrd

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.



Theoretical Prediction of the Heats of Formation, Densities, and Relative Sensitivities for 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo (3:3:0)octan-2-one

by Edward FC Byrd

Weapons and Materials Research Directorate, ARL

REPORT DOCUMENTATION PAGE				<i>Form Approved OMB No. 0704-0188</i>
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.				
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.				
1. REPORT DATE (DD-MM-YYYY) April 2016	2. REPORT TYPE Technical Note	3. DATES COVERED (From - To) 1–31 May 2015		
4. TITLE AND SUBTITLE Theoretical Prediction of the Heats of Formation, Densities, and Relative Sensitivities for 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo (3:3:0)octan-2-one			5a. CONTRACT NUMBER	
			5b. GRANT NUMBER	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Edward FC Byrd			5d. PROJECT NUMBER	
			5e. TASK NUMBER	
			5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Research Laboratory ATTN: RDRL-WML-B Aberdeen Proving Ground, MD 21005-5069			8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TN-0749	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.				
13. SUPPLEMENTARY NOTES				
14. ABSTRACT Using the US Army Research Laboratory-developed series of scripts, written to dramatically simplify the computation of crystalline density and heat of formation, the performance properties for the 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one notional energetic material were evaluated. Additionally, a qualitative estimation of the impact sensitivity has been calculated. This report outlines the procedures used to generate this information, as well as Cheetah calculations using the predicted crystalline density and heat of formation.				
15. SUBJECT TERMS computational toolbox, script, crystalline density, heat of formation, impact sensitivity, energetic materials				
16. SECURITY CLASSIFICATION OF: Unclassified		17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 16	19a. NAME OF RESPONSIBLE PERSON Edward FC Byrd
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	19b. TELEPHONE NUMBER (Include area code) 410-306-0729	

Contents

List of Figures	iv
List of Tables	iv
Acknowledgments	v
1. Introduction	1
2. Results and Discussion	1
3. Conclusions	4
4. References	5
List of Symbols, Abbreviations, and Acronyms	6
Distribution List	7

List of Figures

Fig. 1	Optimized structure of the 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one molecule	1
Fig. 2	Electrostatic potential map of 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one, a) with and b-c) without molecule overlay. Image c) shows the opposite side of the molecule than b)	3

List of Tables

Table 1	Cheetah predicted properties for 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one using predicted density (1.888 g/cm ³).....	4
Table 2	Cheetah predicted properties for 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one using experimental density (1.952 g/cm ³).....	4

Acknowledgments

Dr Betsy Rice and Jennifer J Hare are acknowledged for their efforts in the original coding of the neutral heat of formation and EDAT tools, respectively. Dr Anthony Yau is acknowledged for his work in revising the EDAT code. Dr James Ianni (Applications Engineer with Lockheed-Martin, contractor to the US Army Research Laboratory Department of Defense [DOD] Supercomputing Resource Center [ARL DSRC]) is acknowledged for his “gsubmit” script, initially written for the ARL DSRC. Dr Betsy Rice is acknowledged for running the Cheetah calculations. All computations were performed at the ARL DSRC, Aberdeen Proving Ground, MD. Calculations were performed at the behest of Dr Patrick Caruana (Naval Surface Warfare Center – Indian Head). Drs Michael Chaykovsky and William M Koppes¹ are acknowledged as originators of the molecule modeled herein.

¹ Chaykovsky M, Koppes WM, inventors; The United States of America as represented by the Secretary of the Navy, assignee. 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo[3.3.0]octan-2-one. United States patent US 5,262,544. 1993 Nov 16.

INTENTIONALLY LEFT BLANK.

1. Introduction

US Army Research Laboratory (ARL) researchers have achieved robust theoretical models capable of predicting performance properties, such as heats of formation,^{1,2} densities,^{3,4} and impact sensitivity⁵ of energetic materials and have begun growing advanced synthesis capabilities to realize notional materials. This dual capability allows synthetic and formulation chemists to safely and quickly screen candidate materials to focus efforts only on the most promising compounds. For an in-depth explanation of the different theoretical methods employed herein, please refer to previous works.^{6,7} This technical note will detail theoretical predictions of heat of formation, density, sensitivity, and performance for the 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one molecule.⁸

2. Results and Discussion

The properties of the 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one molecule (Fig. 1) were predicted using the ARL tools.⁷ For the estimation of the impact sensitivities, the electrostatic maps on the 0.001 isosurfaces were generated with the scalar range of the electrostatic surface potential (ESP) ranging from -0.05 to 0.075. Recall that for this visualization methodology, regions of large positive charge (i.e., electron deficient regions, labeled as red) over the backbone of the structure tend to indicate increased sensitivity.




Fig. 1 Optimized structure of the 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one molecule

The computed gas and solid heats of formation using the group additivity method were -82.11 and -106.43 kcal/mol, respectively. The predicted crystalline density was 1.888 g/cm 3 . The experimental density for this compound is known (1.952 g/cm 3), resulting in a -3.3% error, on par with the published root mean square error of approximately 3% .⁴

Additionally, we plot the ESP maps for this molecule (Fig. 2a–c), with and without the molecule overlaid on the ESP. When the images are analyzed, we would quantify the molecule as marginally sensitive, with the large red section being somewhat misleading as it is due to the presence of hydrogens and not the backbone structure. The experimental h_{50} impact sensitivity for this compound is 59 cm (with a reference cyclotrimethylenetrinitramine [RDX] sensitivity of 19 cm). We performed Cheetah 8.0 calculations⁹ to predict the performance parameters using the predicted heat of formation and density. At the Chapman-Jouguet point, Cheetah yields the values shown in Table 1. When the experimental density and the predicted heat of formation are used, Cheetah yields the values shown in Table 2.




Fig. 2 Electrostatic potential map of 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one, a) with and b-c) without molecule overlay. Image c) shows the opposite side of the molecule than b).

Table 1 Cheetah predicted properties for 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one using predicted density (1.888 g/cm³)

Property	Value
Pressure	29.783 GPa
Shock velocity	8.196 km/s
Temperature	2974.1 K
Total energy of detonation	1.109 TNT equivalence (per cm ³) 0.971 TNT equivalence (per gram)

Note: TNT = trinitrotoluene

Table 2 Cheetah predicted properties for 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one using experimental density (1.952 g/cm³)

Property	Value
Pressure	32.491 GPa
Shock velocity	8.454 km/s
Temperature	2876.7 K
Total energy of detonation	1.163 TNT equivalence (per cm ³) 0.986 TNT equivalence (per gram)

3. Conclusions

The ARL-developed software tools were used to predict the heats of formation and crystalline densities of the 5,7-dinitro-5,7-diaza-1,3-dioxabicyclo(3:3:0)octan-2-one molecule. Using this predicted data, we then ran Cheetah calculations to predict the performance of this material. Additionally, we predicted the qualitative impact sensitivity of this compound using electrostatic potential maps. This information has been transitioned back to the requesting synthetic chemist, Dr David Caruana of Naval Surface Warfare Center – Indian Head.

4. References

1. Byrd EFC, Rice BM. Improved prediction of heats of formation of energetic materials using quantum mechanical calculations. *J Phys Chem A.* 2006;110(3):1005–1013; *ibid* 2009;113:5813.
2. Byrd EFC, Rice BM. A comparison of methods to predict solid phase heats of formation of molecular energetic salts. *J Phys Chem A.* 2009;113(1):345–352.
3. Rice BM, Hare JJ, Byrd EFC. Accurate predictions of crystal densities using quantum mechanical molecular volumes. *J Phys Chem A.* 2007;111(42):10874–10879.
4. Rice BM, Byrd EFC. Evaluation of electrostatic descriptors for predicting crystalline density. *J Comp Chem.* 2013;34(25):2146–2151.
5. Rice BM, Hare JJ. A quantum mechanical investigation of the relation between impact sensitivity and the charge distribution in energetic molecules. *J Phys Chem A.* 2002;106(9):1770–1783.
6. Byrd EFC. Theoretical prediction of the heat of formation, density and relative sensitivity for 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine. Aberdeen Proving Ground (MD): Army Research Laboratory (US); 2015 Apr. Report No.: ARL-TN-0680.
7. Byrd EFC. On the failure of correlating partitioned electrostatic surface potentials using bader's atoms-in-molecules theory to impact sensitivities. Aberdeen Proving Ground (MD): Army Research Laboratory (US); 2013 Sep. Report No.: ARL-TR-6206.
8. Caruana P. Naval Surface Warfare Center, Indian Head, MD. Private communication, 2015 May.
9. Bastea S, Fried LE, Glaeseman KR, Howard WM, Kuo IFW, Souers PC, Vitello PA. Cheetah 8.0 thermochemical code. Livermore (CA): Energetic Materials Center: Lawrence Livermore National Laboratory; 2015.

List of Symbols, Abbreviations, and Acronyms

ARL	US Army Research Laboratory
g/cm ³	grams per cubic centimeter
DSRC	Department of Defense (DOD) Supercomputing Resource Center
ESP	electrostatic surface potential
kcal/mol	kilocalories per mole (unit of energy)
RDX	cyclotrimethylenetrinitramine
TNT	trinitrotoluene

1 DEFENSE TECHNICAL
(PDF) INFORMATION CTR
DTIC OCA

2 DIRECTOR
(PDF) US ARMY RESEARCH LAB
RDRL CIO LL
IMAL HRA MAIL & RECORDS
MGMT

1 GOVT PRINTG OFC
(PDF) A MALHOTRA

1 US ARMY ARDEC
(PDF) A DISTASIO

1 US ARMY ARDEC
(PDF) S NICOLICH

17 DIR USARL
(PDF) RDRL WM
B FORCH
J ZABINSKI
RDRL WML
M ZOLTOSKI
RDRL WML A
W OBERLE
RDRL WML B
N TRIVEDI
J MORRIS
B RICE
E BYRD
RDRL WML C
S AUBERT
J SABATINI
J BANNING
RDRL WML D
R BEYER
RDRL WML E
P WEINACHT
RDRL WML F
D LYON
RDRL WML G
W DRYSDALE
RDRL WML H
J NEWILL
RDRL WMP G
R EHLERS

INTENTIONALLY LEFT BLANK.